www.ijarp.com ISSN 2456-9992 Page: 01-19

International Journal Advanced Research Publications

Research Article

Volume: 02
Issue: 10

FORMULATION, EVALUATION, AND EFFICACY OF HERBAL SHAMPOOS: INTEGRATING AYURVEDIC PHARMACOGNOSY WITH MODERN COLLOID AND COSMETIC SCIENCE

Prithviraj Sahebrao Ade[1*], Jui Subhash Sutar[1], Prathmesh Santosh Nimbalkar[1], Jyoti Nilesh jadhavrao[2], Dr. Tushar Shelke[3]

[1]Student, Department of Pharmacy, Genba Sopanrao Moze College of Pharmacy, Wagholi, Pune, Maharashtra, India.

[2]Assistant Professor, Department of Pharmaceutics, Genba Sopanrao Moze College of Pharmacy, Wagholi, Pune, Maharashtra, India.

[3]Principal, Genba Sopanrao Moze College of Pharmacy, Wagholi, Pune, Maharashtra, India.

Article Received: 20 September 2026, Article Revised: 10 October 2026, Published on: 30 October 2026

***Corresponding Author: Prithviraj Sahebrao Ade**

Student, Department of Pharmacy, Genba Sopanrao Moze College of Pharmacy, Wagholi, Pune, Maharashtra, India.

Doi: https://doi-doi.org/101555/ijarp.2703

ABSTRACT

Scalp disorders and hair shaft degradation caused by environmental pollutants, microbial dysbiosis, oxidative stress, and harsh synthetic cleansing agents represent a major global dermatological and cosmetic challenge. Conventional shampoos rely heavily on anionic petrochemical surfactants such as sodium lauryl sulfate (SLS) and sodium laureth sulfate (SLES), alongside synthetic preservatives, silicones, and alkanolamides. While effective at rapid sebum emulsification, prolonged exposure to these synthetic excipients strips the protective 18-methyleicosanoic acid (18-MEA) lipid monolayer, alkalinizes the scalp acid mantle, denatures cuticular alpha-keratin, and triggers corneocyte inflammation, contact dermatitis, and ocular irritation. In recent years, polyherbal shampoos rooted in classical Indian Ayurvedic pharmacognosy (Keshya rasayanas) have attracted significant scientific interest as biocompatible, biodegradable, and multifunctional alternatives. Botanical surfactants rich in triterpenoid and steroidal saponins—principally derived from Sapindus mukorossi (Reetha) and Acacia concinna (Shikakai)—reduce interfacial tension and solubilize sebum via micellar encapsulation without disrupting cuticular integrity. Concurrently, polyphenolic, alkaloid, and mucilage-rich botanicals such as Phyllanthus emblica (Amla), Eclipta prostrata (Bhringraj), Azadirachta indica (Neem), Hibiscus rosa-

sinensis (Japa), Lawsonia inermis (Madayanti), Trigonella foenum-graecum (Methika), Wrightia tinctoria (Indrayava), and Nardostachys jatamansi (Jatamansi) exert potent antioxidant, anti-Malassezia, 5-alpha-reductase inhibitory, and dermal papilla proliferative effects.

KEYWORDS: Herbal Shampoo, Ayurvedic Botanicals, Sapindus mukorossi, Acacia concinna, Saponin Micellar Detergency.

INTRODUCTION

Hair cleansing and scalp hygiene constitute one of the largest segments of the global cosmeceutical and personal care industry, with shampoos serving as the primary pharmaceutical and cosmetic dosage form for managing both aesthetic hair quality and dermatological conditions of the scalp (1,2). Anatomically, the human scalp differs from glabrous skin through its high density of terminal pilosebaceous units (100,000–150,000 follicles), rich neurovascular network, and elevated holocrine sebum secretion rate (1–2 g/day), which collectively create a warm, dark, lipid-rich microenvironment highly susceptible to lipophilic microbial colonization and oxidative lipid peroxidation (3,4). A shampoo is pharmaceutically defined as a viscid aqueous, hydroalcoholic, or solid dispersion of amphiphilic surfactants engineered to remove accumulated sebum, desquamated stratum corneum corneocytes, eccrine sweat residues, environmental particulate matter, and residual hair-styling polymers from the hair shaft and perifollicular ostia without compromising the structural integrity of the hair fiber or the physiological homeostasis of the scalp epidermis (1,5).

Historically, the commercialization of synthetic detergent-based shampoos in the mid-twentieth century replaced traditional soap-based cleansers, which formed insoluble calcium and magnesium carboxylate curds in hard water (5). Contemporary synthetic shampoos predominantly utilize anionic alkyl sulfates and alkyl ether sulfates—most notably sodium lauryl sulfate (SLS) and sodium laureth sulfate (SLES)—at concentrations ranging from 10% to 15% w/w, supplemented with amphoteric foam boosters such as cocamidopropyl betaine (CAPB), nonionic alkanolamide viscosity builders (e.g., cocamide DEA/MEA), synthetic cationic conditioning polymers, polydimethylsiloxanes (dimethicone), parabens, and formaldehyde-releasing preservatives (2,5,6). Although these synthetic matrices produce rapid, voluminous flash foam and efficient lipid stripping, extensive dermatotoxicological and trichological investigations have highlighted their cumulative adverse effects (1,6).

Furthermore, ethoxylated surfactants such as SLES carry the risk of trace contamination with 1,4-dioxane, a probable human carcinogen generated during ethylene oxide ethoxylation, while cocamidopropyl betaine and synthetic preservatives are increasingly implicated in allergic contact dermatitis, scalp pruritus, and disruption of the commensal scalp microbiome (2,6,7).

1. Structural Biology of the Pilosebaceous Unit, Scalp Pathophysiology, and Physicochemical Principles of Detergency




Fig. 1. Schematic representation of the key factors involved in the rational approach to herbal shampoo formulation, illustrating (A) the three-dimensional architecture of the hair shaft, (B) the biochemistry of scalp sebum, and (C) the pathophysiological cascades affecting follicular and cuticular health.

1.1 Ultrastructure and Surface Chemistry of the Hair Shaft

The human hair shaft is a terminally differentiated, keratinized cylindrical filament composed of three concentric morphological zones: the outermost cuticle, the central cortex, and the innermost medulla (4). The cuticle consists of 6 to 10 overlapping layers of flattened, anucleate corneocytes (scales) oriented toward the distal tip, each measuring approximately 0.3–0.5 µm in thickness (4,6). Each cuticular cell is subdivided into the A-layer (high cystine content, >30%), the exocuticle (~15% cystine), the endocuticle (low cystine, ~3%, hydrophilic and swellable), and the cell membrane complex (CMC), which acts as a continuous intercellular lipid-protein adhesive binding adjacent cuticular and cortical cells (4). Crucially, the outermost epicuticle is covered by a monomolecular layer of 18-methyleicosanoic acid (18-MEA) covalently linked to underlying cysteine residues via

thioester bonds (4,6). This 18-MEA lipid envelope imparts natural hydrophobicity, low surface friction, and aesthetic gloss to virgin hair.

1.2 Sebum Biochemistry and the Scalp Acid Mantle

The scalp harbours roughly 100,000 to 150,000 terminal hair follicles associated with multilobular holocrine sebaceous glands that secrete 1–2 g of sebum daily onto the infundibular ostia and proximal hair shaft (3,14). Native human sebum is a complex hydrophobic lipid mixture comprising triglycerides and free fatty acids (57.5%), wax esters (26.0%), squalene (12.0%), and cholesterol/cholesterol esters (4.5%) (3,15). Concurrently, eccrine sweat glands secrete water, lactic acid, urea, pyrrolidone carboxylic acid (PCA), and electrolytes. The emulsification of sebum with eccrine secretions forms the hydrolipidic acid mantle of the scalp, maintaining a physiological pH between 4.5 and 5.5 (4,14). This mildly acidic milieu suppresses pathogenic bacterial proliferation, stabilizes epidermal lipid lamellae synthesis (ceramides via beta-glucocerebrosidase and acidic sphingomyelinase), and keeps the hair cuticle scales tightly appressed (4,14).

1.3 Pathophysiology of Scalp Disorders and Surfactant-Induced Hair Damage

Disruption of scalp homeostasis by aggressive synthetic surfactants, ultraviolet radiation, atmospheric particulate matter, or microbial dysbiosis initiates a cascade of structural and inflammatory pathologies (Table 1) (3,6,15,16).

Table 1 Pathophysiological comparison of synthetic detergent-induced scalp and hair damage versus the therapeutic modulation achieved by Ayurvedic polyherbal shampoos.

Pathophysiological / Biophysical Parameter	Synthetic Sulfate-Based Shampoos	Ayurvedic Polyherbal Shampoos	Underlying Molecular & Biophysical Mechanism	Ref.
Cuticular 18-MEA Lipid Layer	Progressive hydrolytic and surfactant-mediated stripping of covalently bound 18-MEA	Preservation of 18-MEA monolayer; deposition of protective botanical mucilage film	Nonionic/weakly acidic triterpenoid saponins exhibit low keratin-binding affinity compared to strongly anionic -SO4- headgroups	(4,6,10)
Scalp Acid Mantle & Keratin Charge	Often neutral to alkaline (pH 6.5–8.0); increases net negative charge above pI (3.67), causing scale lifting	Naturally acidic (pH 4.5–5.6) due to organic acids (ascorbic, gallic, tartaric); maintains isoionic point (~5.6)	Acidic vehicle protonates basic amino acid residues, minimizing electrostatic repulsion between cuticular scales	(4,9,13,14)

1.4 Colloid and Interfacial Physics of Herbal Saponin Detergency




Fig. 2. Structural comparison of synthetic sulfate surfactants and plant-derived saponins.

When dissolved in an aqueous shampoo vehicle, saponin monomers adsorb at the air–water and oil–water interfaces, orienting their lipophilic triterpenoid/steroidal rings toward the hydrophobic sebum phase (or air) and extending their hydrated sugar chains into the bulk aqueous phase (17). The surface excess concentration (Γ) and interfacial adsorption of saponins are governed by the Gibbs Adsorption Isotherm: $\Gamma = -(1 / RT) \cdot (d\gamma / d \ln C)$, where γ is the surface tension (mN/m), C is the bulk surfactant concentration, R is the universal gas constant, and T is the absolute temperature (17). As the bulk concentration exceeds the Critical Micelle Concentration (CMC)—which for Sapindus mukorossi saponins ranges between 0.015% and 0.045% w/v (nearly an order of magnitude lower than the ~0.23% w/v CMC of SLS)—saponin monomers spontaneously self-assemble into spherical, rod-like, or vesicular micelles characterized by a negative standard free energy of micellization ($\Delta G°mic = RT \ln X_CMC$) (10,17).

2. Pharmacognosy, Phytochemistry, and Molecular Mechanisms of Ayurvedic Hair-Care Botanicals

Table 2. Primary natural surfactants and cleansers used in Ayurvedic shampoo formulations.

Botanical source	Part used	Major bioactive constituents (natural surfactants)	Key functions	References
Soapnut (*Sapindus mukorossi*)	Pericarp	Saponins (sapindosides A and B)	• Natural cleansing agent • Foaming and emulsifying • Removes excess sebum dirt and pollutants	[9,10]
Shikakai (*Acacia concinna*)	Pods	Triterpenoid saponins (acacic acid, concinnosides)	• Mild cleanser • Maintains hair softness • Reduces static charge	[11,12]
Reetha (*Sapindus trifoliatus*)	Pericarp	Saponins (trifoliatosides)	• Effective surfactant • Removes oil and impurities • Improves foam stability	[13,14]
Amla (*Phyllanthus emblica*)	Fruit	Saponins, polyphenols	• Gentle cleansing • Antioxidant support	[15,16]
Fenugreek (*Trigonella foenum-graecum*)	Seeds	Saponins (diosgenin derivatives)	• Mild surfactant • Conditioning effect	[17,18]

Table 3. Secondary botanicals used for conditioning, antioxidant, anti-dandruff and follicular-stimulating activities in Ayurvedic shampoos.

Botanical source	Part used	Major bioactive constituents	Key functions	References
Aloe vera (*Aloe barbadensis*)	Leaf gel	Polysaccharides (acemannan), vitamins, minerals	• Conditioning • Moisturizing • Soothes scalp irritation	[19,20]
Bhringraj (*Eclipta prostrata*)	Whole plant	Wedelolactone, flavonoids	• Promotes hair growth • Anti-inflammatory	[21,22]
Neem (*Azadirachta indica*)	Leaves	Nimbidin, azadirachtin, flavonoids	• Anti-dandruff • Antimicrobial • Controls *Malassezia* spp.	[23,24]
Hibiscus (*Hibiscus rosa-sinensis*)	Flower	Flavonoids, anthocyanins, organic acids	• Conditioning • Strengthens hair shaft • Improves shine	[25,26]
Green tea (*Camellia sinensis*)	Leaves	Catechins (EGCG), polyphenols	• Antioxidant • Inhibits DHT • Supports hair follicle health	[27,28]
Licorice (*Glycyrrhiza glabra*)	Root	Glycyrrhizin, flavonoids	• Anti-inflammatory • Soothes scalp • Reduces irritation	[29,30]




Fig. 3. Multifaceted mechanisms of Ayurvedic botanicals in the pilosebaceous unit. Primary natural surfactants cleanse and regulate sebum, while secondary botanicals exhibit conditioning, antioxidant, anti-dandruff and follicular-stimulating activities through modulation of multiple molecular signaling pathways.

Table 2 Pharmacognostic profile, phytochemical markers, and cosmeceutical functions of primary cleansing saponin botanicals used in Indian herbal shampoos.

Botanical Name, Family & Ayurvedic Name	Plant Part Used & API Yield Standard	Principal Phytochemical Markers	Functional Role Shampoo Formulation	Key Pharmacological	Ref.
Sapindus mukorossi Gaertn. Arishtaka / Reetha	Dried fruit pericarp (Total saponins: 10.1–11.5% w/w)	Aglycone: Hederagenin (C30 oleanane) Glycosides: Sapindosides A–E, Mukorozisaponins E1, G, Y1, Y2; Mukuroziosides Ia–IIb	Primary surfactant, foaming agent, wetting agent, and O/W sebum emulsifier	Lowers γ_LV to ~33 mN/m; inhibits Malassezia furfur (MIC 62.5 µg/mL); downregulates COX-2 and 5-LOX in scalp keratinocytes	(9,10,18)
Acacia concinna (Willd.) DC. Saptala / Shikakai	Dried pods and bark (Total saponins: 11.0–16.0% w/w)	Aglycone: Acacic acid (C30 oleanane) Glycosides: Kinmoonosides A–C, Acaciasides A & B; Tartaric, citric & ascorbic acids; Lupeol	Mild co-surfactant, natural pH acidifier (pH 4.5–5.2), detangler, and cuticular conditioner	Seals raised cuticular scales; reduces wet combing force; exhibits antifungal activity against Trichophyton and Malassezia spp.	(7,9,19)
Ziziphus jujuba Mill. / Z. spina-christi Badara / Ber / Sidr	Dried leaves (Saponins: 3.5–6.0%; Mucilage: 8–12%)	Aglycone: Jujubogenin (dammarane) Glycosides: Ziziphin, Jujubosides A & B, Christinin-A; Rhamnogalacturonan mucilage	Foam stabilizer, natural viscosity modifier, and hair shaft film-former	Retards plateau-border liquid drainage in foam lamellae; provides smooth sensory slip and antioxidant protection	(1,13,20)

2.1 Sapindus mukorossi Gaertn. and Sapindus trifoliatus L. (Sapindaceae) — Reetha / Arishtaka

The dried fruit pericarp of Sapindus mukorossi (predominantly North Indian and Himalayan distribution) and Sapindus trifoliatus / S. laurifolius (South and Western Indian distribution), known in Sanskrit as Arishtaka or Phenila ('foam-producing'), represents the cornerstone primary surfactant of Indian herbal shampoos (9,10,18). The pericarp contains 10.1% to 11.5% w/w (and up to 28% in methanolic extracts) of triterpenoid saponins classified into three structural subclasses based on their aglycone skeleton: (i) Oleanane-type monodesmosidic and bisdesmosidic saponins (Sapindosides A–E and Mukorozisaponins E1, G, Y1, Y2), which possess hederagenin (3β,23-dihydroxyolean-12-en-28-oic acid) as the primary aglycone linked to L-arabinose, L-rhamnose, and D-xylose moieties; (ii) Dammarane-type tetracyclic triterpenoid saponins (Sapimukosides A–J); and (iii) Tirucallane-type triterpenoid saponins (Sapinmusaponins A–E) (18). Crucially, the pericarp also yields acyclic sesquiterpene oligoglycosides (Mukuroziosides Ia, Ib, IIa, IIb), which act as natural hydrotropes and co-surfactants that enhance the aqueous solubility and wetting kinetics of the hederagenin saponins (18).

2.2 Acacia concinna (Willd.) DC. [syn. Senegalia rugata] (Fabaceae) — Shikakai / Saptala

Known classically as Saptala or Shikakai ('fruit for the hair' in Tamil), the dried pods of the prickly climbing shrub Acacia concinna serve as the primary mild co-surfactant, detangler, and cuticular conditioner in traditional and commercial Indian shampoos (1,7,9). The pods and bark of A. concinna contain 11.0% to 16.0% w/w of acylated pentacyclic triterpenoid saponins built around the aglycone acacic acid (3β,16α,21β-trihydroxyolean-12-en-28-oic acid) (19). The major characterized bioactive macromolecular saponins include Kinmoonosides A, B, and C, and Acaciasides A and B, wherein the acacic acid sapogenin is esterified at C-21 with monoterpene carboxylic acids [specifically (2E,6S)-2,6-dimethyl-6-hydroxyocta-2,7-dienoic acid] and glycosylated at C-3 and C-28 with oligosaccharide chains containing D-glucose, L-arabinose, L-rhamnose, and D-xylose (19).

2.3 Ziziphus jujuba Mill. / Z. spina-christi (Rhamnaceae) and Albizia amara (Roxb.) Boivin (Fabaceae) — Ber & Arappu

In regional Indian formulations (particularly South Indian Arappu and North/Western Indian Badara preparations), the pulverized leaves of Ziziphus jujuba (and Z. spina-christi) and Albizia amara are co-formulated with Sapindus and Acacia to stabilize foam elasticity and increase dispersion viscosity (1,13,20,21). Ziziphus leaves contain dammarane-type

saponins—notably Ziziphin, Jujubosides A and B, and Christinin-A (derived from the aglycone jujubogenin)—alongside high concentrations of mucilage polysaccharides (rhamnogalacturonans and arabinogalactans), rutin, and quercetin-3-O-robinobioside (13,20). Albizia amara leaves yield oleanane saponins (echinocystic acid and machaerinic acid glycosides) and macrocyclic spermine alkaloids (budmunchiamines A–C) (7,21). In a polyherbal shampoo system, the high-molecular-weight mucilage polysaccharides of Ziziphus and Albizia migrate to the plateau borders (lamellae) between saponin foam bubbles, increasing bulk liquid viscosity and retarding gravitational drainage of water from the foam film (13,17,21).

2.4 Secondary Conditioning, Anti-Dandruff, and Follicular-Stimulating Indian Botanicals

To transform a basic saponin cleanser into a therapeutic cosmeceutical shampoo, classical Ayurvedic formulations incorporate secondary botanicals that target dermal papilla mitogenesis, melanogenesis, cuticular conditioning, and microbial dysbiosis (Table 3) (7,8,11,12).

Table 3 Secondary conditioning, anti-dandruff, and follicular-stimulating Indian botanicals, their standardization markers, and molecular mechanisms.

Botanical Name, Family & Ayurvedic Name	Part Used & Standardization Marker	Key Bioactive Phytoconstituents	Molecular & Cellular Targets in Scalp/Follicle	Cosmeceutical & Clinical Benefit in Shampoo	Ref.
Phyllanthus emblica L. Amalaki / Amla	Dried fruit pericarp (Gallic acid ≥1.5% w/w; Ascorbic acid; Tannins ≥25%)	Emblicanin A & B, Punigluconin, Pedunculagin, Gallic acid, Ellagic acid, Corilagin, Chebulagic acid	Inhibits 5α-reductase type II; activates Nrf2/HO-1 antioxidant axis; chelates pro-oxidant Fe2+/Cu2+; stimulates DP proliferation	Prevents squalene peroxidation and premature greying (canities); astringes cuticular scales to boost specular gloss	(7,23,24)
Eclipta prostrata (L.) L. [syn. E. alba]	Whole aerial plant (Wedelolactone ≥0.5–1.2% w/w)	Coumestans (Wedelolactone, Demethylwedelolactone), Eclalbasaponins I–	Downregulates TGF-β1 and DKK-1; upregulates Wnt/β-	Accelerates telogen-to-anagen transition; increases	(7,11,25)

(Asteraceae) Bhringraj / Kesharaja		VI, Luteolin, Apigenin, β-amyrin	catenin, Shh, FGF-7, and VEGF in dermal papilla cells; activates follicular melanocytes	hair follicle length and bulb diameter; restores melanin synthesis	
Azadirachta indica A. Juss. (Meliaceae) Nimba / Neem	Dried leaves & seed oil (Azadirachtin ≥0.1%; Nimbidin ≥1.0%)	Limonoids/Triterpenoids (Azadirachtin, Nimbin, Nimbidin, Salannin, Gedunin), Quercetin, Mahmoodin	Disrupts Malassezia furfur cell membrane and inhibits extracellular fungal lipase; blocks NF-κB p65 nuclear translocation and COX-2	Potent anti-dandruff and antipruritic agent; resolves seborrheic dermatitis, scalp folliculitis, and pediculosis	(15,26,27)
Hibiscus rosa-sinensis L. (Malvaceae) Japa / Jaswand	Flowers and leaves (Total anthocyanins ≥0.8%; Mucilage ≥12%)	Cyanidin-3-sophoroside, Cyanidin-3,5-diglucoside, Quercetin-3-diglucoside, L-rhamnose/D-galacturonic acid mucilage	Upregulates VEGF and IGF-1 in perifollicular vasculature; acidic mucilage forms hydrated polyelectrolyte film on keratin	Natural cationic-mimetic detangler; reduces wet combing force by 35–45%; prevents trichoptilosis (split ends)	(7,12,28)

2.5 Extraction Standardization, Green Extraction Technologies, and Phyto-Synergism

The therapeutic and surface-active reproducibility of an herbal shampoo depends critically on extraction methodology (10,17). Traditional open-vessel boiling (Kwatha decoction at 100°C for 2–4 hours) causes hydrolytic cleavage of C-28 ester glycosidic linkages in bisdesmosidic saponins (Sapindosides and Acaciasides), converting them into less water-soluble monodesmosides and lowering foaming capacity, while thermally degrading heat-labile ascorbic acid and anthocyanins (17,18,23). Modern pharmaceutical manufacturing employs Ultrasound-Assisted Extraction (UAE, 20–40 kHz at 45°C using 50–70% v/v hydroethanolic menstruum) or Microwave-Assisted Extraction (MAE), which induce acoustic cavitation and

rapid intracellular pressure buildup to rupture lignified pericarp cell walls within 20–30 minutes, increasing total saponin yield by 28–36% while preserving polyphenolic integrity (17,18). Furthermore, spray-drying of standardized hydroalcoholic extracts with 5% maltodextrin yields free-flowing, non-hygroscopic phyto-powders standardized by High-Performance Thin-Layer Chromatography (HPTLC) or RP-HPLC against reference markers: Hederagenin (≥2.5% w/w in Sapindus extract), Acacic acid lactone (≥2.0% w/w in Acacia extract), Gallic acid (≥3.0% w/w in Phyllanthus extract), and Wedelolactone (≥0.8% w/w in Eclipta extract) (18,19,23,25).

3. Formulation Development and Pharmaceutical Engineering of Herbal Shampoos

Formulating a stable, aesthetically acceptable, and high-performing herbal shampoo requires balancing opposing colloid and rheological forces (1,5,10). While synthetic shampoos rely on simple ternary surfactant–salt–water phase diagrams (where adding 1–2% NaCl induces a sphere-to-wormlike micellar transition in SLES/CAPB systems, effortlessly raising viscosity to 4,000–8,000 cP), natural triterpenoid saponins do not form entangled wormlike micelles upon NaCl addition (5,10,17). Therefore, herbal shampoo engineering demands specialized surfactant blending, hydrocolloid network structuring, chelation, pH buffering, and green preservation (Table 4 & Fig. 4).

3.1 Purely Herbal (100% Botanical) vs. Eco-Certified Green Surfactant Hybrid Matrices

In contemporary cosmetic pharmaceutics, herbal shampoos are engineered into two distinct architectures (Table 4) (2,5,10,13):

Table 4 Master pharmaceutical formulation engineering matrices comparing (A) 100% Purely Botanical Ayurvedic Shampoo, (B) Eco-Certified Green Surfactant Hybrid Shampoo, and (C) Waterless Solid Polyherbal Syndet Bar.

Functional Excipient Category	Formulation A: 100% Purely Botanical Ayurvedic Shampoo	Formulation B: Eco-Certified Green Surfactant Hybrid Herbal Shampoo	Formulation C: Waterless Solid Polyherbal Syndet Shampoo Bar	Specific Pharmaceutical & Colloid Engineering Role	Ref.
Primary Cleansing Surfactant System	Sapindus mukorossi standardized pericarp	Sapindus mukorossi extract: 6.0% Coco-Glucoside	Sodium Cocoyl Isethionate (SCI): 48.0%	Reduces γ_LV to 32–35 mN/m; initiates roll-up emulsification	(5,10,35)

	extract: 12.0% Acacia concinna pod extract: 6.0%	(APG): 8.0% Sodium Cocoyl Glutamate: 5.0%	Coco-Glucoside: 10.0% Sapindus spray-dried powder: 10.0%	of sebum; generates stable viscoelastic foam	
Co-Surfactant & Foam Booster	Ziziphus jujuba leaf extract: 3.0% Albizia amara extract: 2.0%	Acacia concinna extract: 3.0% Cocamidopropyl Betaine (plant-derived): 4.0%	Acacia concinna pod powder: 8.0% Stearic Acid / Cetyl Alcohol (binder): 8.0%	Packs at air–water interface via H-bonding; increases surface dilational modulus and flash foam volume	(5,13,21)
Active Therapeutic Keshya Botanicals	Phyllanthus emblica: 2.0% Eclipta prostrata: 2.0% Azadirachta indica: 1.5% Lawsonia inermis: 1.0%	Phyllanthus emblica: 1.5% Eclipta prostrata: 1.5% Azadirachta indica: 1.0% Wrightia tinctoria: 1.0%	Phyllanthus emblica powder: 3.0% Eclipta prostrata extract: 2.0% Azadirachta seed oil: 1.5% Hibiscus flower powder: 2.5%	Delivers 5α-reductase inhibition, Wnt/β-catenin activation, anti-Malassezia action, and cuticular scale sealing	(7,11,23,26)

3.2 Rheological Engineering and Non-Newtonian Pseudoplasticity

An ideal liquid herbal shampoo must exhibit non-Newtonian pseudoplastic (shear-thinning) rheology governed by the Ostwald–de Waele Power Law Equation: $\tau = K \cdot \dot{\gamma}^n$, and apparent viscosity $\eta = K \cdot \dot{\gamma}^{(n-1)}$, where τ is shear stress (Pa), $\dot{\gamma}$ is shear rate (s^{-1}), K is the consistency index, and n is the flow behaviour index ($n < 1$ for pseudoplastic fluids) (10,13,36). At rest inside the bottle (low shear rate, $\dot{\gamma} < 0.1\ s^{-1}$), the shampoo must maintain a high apparent viscosity (2,500–5,500 cP at 25°C) and yield stress to prevent gravitational sedimentation of colloidal herbal particulates and phase separation of essential oils (10,36). Conversely, during squeezing from the orifice and manual spreading across wet hair (high shear rate, $\dot{\gamma} = 50–500\ s^{-1}$), the apparent viscosity must drop rapidly (to 300–700 cP) to allow effortless dispensing and uniform lather distribution (Fig. 4A) (36).

3.3 Hard-Water Compatibility, Chelating Systems, and Isoionic pH Buffering

A classic failure mode of crude Ayurvedic decoctions in Indian household settings—where municipal and groundwater hardness frequently ranges from 150 to 400 ppm CaCO3 equivalent—is the complexation of divalent cations (Ca2+ and Mg2+) with the C-28 carboxylic acid groups (-COO^-) of glucuronic acid-bearing saponins and hydrolysable gallotannins, resulting in foam collapse and dulling mineral deposits on the hair shaft (9,10). Furthermore, trace transition metal ions (Fe2+, Fe3+, Cu2+) leached from plant raw materials catalyze the auto-oxidation of gallic acid, catechin, and wedelolactone into dark ortho-quinones, causing severe colour darkening and sediment formation during shelf storage (23).

3.4 Green Preservation Strategies for Carbohydrate-Rich Botanical Matrices

Because polyherbal shampoos contain high levels of botanical sugars, free oligosaccharides, amino acids, and mucilage hydrocolloids, they provide an ideal nutrient broth for Gram-negative bacteria (especially Pseudomonas aeruginosa and Burkholderia cepacia), Gram-positive cocci (Staphylococcus aureus), yeasts (Candida albicans), and moulds (Aspergillus brasiliensis) (10,37). Moreover, high concentrations of nonionic saponins and ethoxylated/glucoside surfactants can partition lipophilic preservatives (such as parabens or phenoxyethanol) into the hydrophobic core of surfactant micelles, drastically reducing the unbound preservative concentration in the bulk aqueous phase where microbes reside (37).

4. Physicochemical, Analytical, Microbiological, and Clinical Evaluation Protocols

Rigorous quality control of herbal shampoos requires standardized evaluation protocols encompassing physicochemical performance, rheological stability, phytochemical marker quantification, microbiological safety, and clinical trichological efficacy, benchmarked against the Bureau of Indian Standards (BIS) specification IS 7884:2023 (and IS 7884:2004 reaffirmations), AYUSH Schedule T Good Manufacturing Practices, and the Ayurvedic Pharmacopoeia of India (API) (Table 5) (9,10,13,38,39).

Table 5 Standardized physicochemical, analytical, microbiological, and toxicological evaluation protocols and regulatory acceptance benchmarks (BIS IS 7884:2023 & AYUSH API).

Evaluation Parameter	Standardized Experimental /Protocol	Mathematical Instrumentation	BIS IS Benchmark	Ref.
pH Determination	Potentiometric measurement of 10% w/v shampoo dispersion in freshly boiled and cooled distilled water at 25 ± 1°C	Calibrated digital glass-electrode pH meter	BIS IS 7884 range: 4.0–9.0 Optimal Trichological Target: 4.8–5.6	(9,14,38)
Percent Solid Content (% w/w)	Gravimetric evaporation of 4.0 g shampoo in a tared porcelain evaporating dish on a water bath followed by hot-air oven drying at 105 ± 2°C to constant weight	% Solids = (W_dry / W_initial) × 100	20.0% – 30.0% w/w (Ensures easy washability without sticky residue)	(9,13,38)
Surface Tension Reduction (γ_LV)	Drop-weight/drop-count method using a Traube Stalagmometer or Du Noüy ring tensiometer on a 10% w/v aqueous shampoo dilution at 25°C	$\gamma_2 = \gamma_1 \cdot (n_1 \cdot \rho_2) / (n_2 \cdot \rho_1)$ (where γ_1 of water = 72.8 mN/m)	31.0 – 38.0 mN/m (dynes/cm) (Must reduce water γ by >50%)	(9,10,13)
Foaming Ability & Stability	Ross–Miles Foam Analyzer (ASTM D1173) or Cylinder Shake Method (50 mL of 1% w/v shampoo shaken 10 times; foam volume recorded at t = 1 min and t = 5 min)	Foam Stability Index (FSI) = (V_5min / V_1min) × 100	BIS IS 7884 Min Foam Height: ≥150 mm at 5 min FSI: ≥85% retention over 5 min	(9,13,38)
Wetting Time	Draves–Clarkson / Canvas Disc Method: 1-inch diameter standard canvas disc (0.44 g) floated on 1% w/v shampoo surface; time to start sinking recorded via stopwatch	Direct chronometric measurement (seconds) at 25°C	110 – 185 seconds for purely herbal; 65 – 120 seconds for green hybrid shampoos	(9,10,13)
Detergency & Sebum Cleaning Power	Barnett and Powers Test: 5.0 g human hair swatch soiled with artificial sebum (squalene, olive oil, coconut oil, oleic acid, cholesterol) in hexane; washed in 100 mL of 5% shampoo at 35°C	DP (%) = [1 - (T / C)] × 100 (T = residual sebum in test; C = initial sebum in control)	Purely Botanical: 58% – 72% Green Hybrid Herbal: 74% – 86%	(9,10,13,40)

5. Future Aspects: Phyto-Nanotechnology, Network Pharmacology, and Sustainable Upcycling

A fundamental biopharmaceutical limitation of conventional herbal shampoos is their short contact residence time on the scalp (typically 1.5 to 3.0 minutes prior to aqueous rinsing) coupled with the poor stratum corneum permeability of polar, high-molecular-weight polyphenolic glycosides (such as wedelolactone, corilagin, and emblicanins, which violate Lipinski's Rule of Five due to high topological polar surface area and multiple hydrogen-bond donors) (23,42). Consequently, over 95% of unencapsulated water-soluble herbal bioactives are washed down the drain before reaching the follicular bulge or dermal papilla (42,43). Integrating phyto-nanotechnology, network pharmacology, and circular green chemistry represents the transformative frontier of herbal trichology

CONCLUSION

The resurgence of Ayurvedic herbal shampoos represents far more than a nostalgic return to traditional ethnobotanical remedies; when engineered through the lens of modern colloid chemistry, trichological biophysics, and pharmaceutical technology, polyherbal shampoos offer a scientifically superior, multi-targeted alternative to harsh petrochemical sulfate detergents. While synthetic SLS and SLES shampoos strip the protective 18-MEA cuticular lipid monolayer, alkalinize the scalp acid mantle, and perturb the epidermal barrier, triterpenoid and steroidal saponins from Sapindus mukorossi, Acacia concinna, Ziziphus jujuba, and Albizia amara gently solubilize excess sebum via micellar roll-up at an isoionic pH (4.8–5.6) that keeps cuticular scales tightly appressed. Simultaneously, secondary Keshya and Kandughna botanicals—including Phyllanthus emblica, Eclipta prostrata, Azadirachta indica, Hibiscus rosa-sinensis, Lawsonia inermis, Trigonella foenum-graecum, and Wrightia tinctoria—intervene across the molecular pathophysiological network of scalp disorders by inhibiting 5α-reductase, activating canonical Wnt/β-catenin signaling in dermal papilla cells, complexing fungal ergosterol in Malassezia spp., scavenging squalene peroxides, and cross-linking damaged keratin fibers. Overcoming the historical limitations of crude herbal decoctions requires rigorous standardization of marker phytoconstituents (via HPTLC/HPLC), synergistic blending with eco-certified alkyl polyglucosides and hydrocolloid networks (xanthan–galactomannan), green chelation with sodium phytate, and incorporation of 200–600 nm phyto-nanocarriers (phytosomes, niosomes, transfersomes, and NLCs) that leverage the cuticular ratchet effect to establish sustained follicular depots during brief wash cycles. Continued integration of network pharmacology, BIS IS 7884:2023 quality

validation, and randomized phototrichogram clinical trials will be pivotal in translating classical Indian pharmacognosy into globally competitive, evidence-based cosmeceutical hair therapeutics.

ACKNOWLEDGEMENTS

The authors would like to express their sincere gratitude to **Ms. Jyoti Nilesh jadhavrao** Assistant Professor, Department of Pharmaceutics for their valuable guidance, constructive suggestions, and continuous support throughout the preparation of this review article. The authors also gratefully acknowledge **Dr. Tushar Shelke** Principal, for providing the necessary institutional support, encouragement, and facilities required for the completion of this work. The authors sincerely appreciate their valuable contributions and support.

REFERENCES

1. Arora P, Nanda A, Karan M. Shampoos based on synthetic ingredients vis-a-vis shampoos based on herbal ingredients: a review. Int J Pharm Sci Rev Res. 2011 Mar-Apr;7(1):41-6. Available from: https://globalresearchonline.net/journalcontents/volume7issue1/article-007.pdf
2. Gubitosa J, Rizzi V, Fini P, Cosma P. Hair care cosmetics: from traditional shampoo to solid/waterless formulations. A review. Cosmetics. 2019 Mar;6(1):13. Available from: https://doi.org/10.3390/cosmetics6010013
3. Draelos ZD. Essentials of hair care often neglected: hair cleansing. Int J Trichology. 2010 Jan;2(1):24-9. Available from: https://doi.org/10.4103/0974-7753.66909
4. Gavazzoni Dias MF. Hair cosmetics: an overview. Int J Trichology. 2015 Jan-Mar;7(1):2-15. Available from: https://doi.org/10.4103/0974-7753.153450
5. Cornwell PA. A review of shampoo surfactant technology: consumer benefits, raw materials and recent developments. Int J Cosmet Sci. 2018 Feb;40(1):16-30. Available from: https://doi.org/10.1111/ics.12439
6. Robbins CR. Chemical and physical behavior of human hair. 5th ed. Berlin: Springer-Verlag; 2012. Chapter 5, Interactions of shampoo and crème rinse ingredients with human hair; p. 329-443. Available from: https://doi.org/10.1007/978-3-642-25611-0_5
7. Patel S, Sharma V, Chauhan NS, Thakur M, Dixit VK. Hair growth: focus on herbal therapeutic agent. Curr Drug Discov Technol. 2015;12(1):21-42. Available from: https://doi.org/10.2174/1570163812666150610115055
8. Hazra J, Panda AK. Concept of Keshya drugs (hair promoter) in Ayurveda. Int J Res Ayurveda Pharm. 2013;4(1):6-9. Available from: https://doi.org/10.7897/2277-4343.04112

9. Mainkar AR, Jolly CI. Evaluation of commercial herbal shampoos. Int J Cosmet Sci. 2000 Oct;22(5):385-91. Available from: https://doi.org/10.1046/j.1467-2494.2000.00047.x
10. Mainkar AR, Jolly CI. Formulation of natural shampoos. Int J Cosmet Sci. 2001 Feb;23(1):59-62. Available from: https://doi.org/10.1046/j.1467-2494.2001.00057.x
11. Roy RK, Thakur M, Dixit VK. Hair growth promoting activity of Eclipta alba in male albino rats. Arch Dermatol Res. 2008 Aug;300(7):357-64. Available from: https://doi.org/10.1007/s00403-008-0860-3
12. Adhirajan N, Ravi Kumar T, Shanmugasundaram N, Babu M. In vivo and in vitro evaluation of hair growth potential of Hibiscus rosa-sinensis Linn. J Ethnopharmacol. 2003 Oct;88(2-3):235-9. Available from: https://doi.org/10.1016/S0378-8741(03)00231-9
13. Al Badi K, Khan SA. Formulation, evaluation and comparison of the herbal shampoo with the commercial shampoos. Beni-Suef Univ J Basic Appl Sci. 2014 Dec;3(4):301-5. Available from: https://doi.org/10.1016/j.bjbas.2014.11.005
14. Gavazzoni Dias MF, de Almeida AM, Cecato PM, Adriano AR, Pichler J. The shampoo pH can affect the hair: myth or reality? Int J Trichology. 2014 Jul;6(3):95-9. Available from: https://doi.org/10.4103/0974-7753.139078
15. Ro BI, Dawson TL. The role of sebaceous gland activity and scalp microfloral metabolism in the etiology of seborrheic dermatitis and dandruff. J Investig Dermatol Symp Proc. 2005 Dec;10(3):194-7. Available from: https://doi.org/10.1111/j.1087-0024.2005.10104.x
16. Trüeb RM. Oxidative stress in ageing of hair. Int J Trichology. 2009 Jan;1(1):6-14. Available from: https://doi.org/10.4103/0974-7753.51923
17. Güçlü-Üstündağ Ö, Mazza G. Saponins: properties, applications and processing. Crit Rev Food Sci Nutr. 2007;47(3):231-58. Available from: https://doi.org/10.1080/10408390600698197
18. Suhagia BN, Rathod IS, Sindhu S. Sapindus mukorossi (Areetha): an overview. Int J Pharm Sci Res. 2011;2(8):1905-13. Available from: https://doi.org/10.13040/IJPSR.0975-8232.2(8).1905-13
19. Tezuka Y, Honda K, Banskota AH, Thet MM, Kadota S. Kinmoonosides A–C, three new cytotoxic saponins from the fruits of Acacia concinna, a medicinal plant collected in Myanmar. J Nat Prod. 2000 Dec;63(12):1658-64. Available from: https://doi.org/10.1021/np000347f
20. Aghel N, Moghimipour E, Dana AR. Formulation of a herbal shampoo using total saponins of Acanthophyllum squarrosum. Iran J Pharm Res. 2007;6(3):167-72. Available from: https://pubmed.ncbi.nlm.nih.gov/24250504/

21. Indalkar Y, Pimpodkar N, Godase A, Gaikwad P. A compressive review on Albizia amara: a potential plant-derived surfactant for cosmetic and pharmaceutical applications. J Pharmacogn Phytochem. 2025;14(1):112-20. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC12786934/
22. Wani SA, Kumar P. Fenugreek: a review on its nutraceutical properties and utilization in various food and cosmetic products. J Saudi Soc Agric Sci. 2018 Apr;17(2):97-106. Available from: https://doi.org/10.1016/j.jssas.2016.01.007
23. Luanpitpong S, Nimmannit U, Pongrakhananon V, Chanvorachote P. Emblica (Phyllanthus emblica Linn.) fruit extract promotes proliferation in dermal papilla cells of human hair follicle. Res J Med Plant. 2011;5(1):95-100. Available from: https://doi.org/10.3923/rjmp.2011.95.100
24. Majeed M, Majeed S, Nagabhushanam K, Mundkur L, Neupane P, Shah K. Clinical study to evaluate the efficacy and safety of a hair serum product in healthy adult male and female volunteers with hair fall. Clin Cosmet Investig Dermatol. 2020 Sep 24;13:691-700. Available from: https://doi.org/10.2147/CCID.S271013
25. Datta K, Singh AT, Mukherjee A, Bhat B, Ramesh B, Burman AC. Eclipta alba extract with potential for hair growth promoting activity. J Ethnopharmacol. 2009 Jul 30;124(3):450-6. Available from: https://doi.org/10.1016/j.jep.2009.05.023
26. Anand N, Aquicio JM, Anand A. Antifungal properties of Neem (Azadirachta indica) leaves extract to treat hair dandruff. E-Int Sci Res J. 2010;2(3):244-52. Available from: https://doi.org/10.37022/jis.v8i3.122
27. Alzohairy MA. Therapeutics role of Azadirachta indica (Neem) and their active constituents in diseases prevention and treatment. Evid Based Complement Alternat Med. 2016;2016:7382506. Available from: https://doi.org/10.1155/2016/7382506
28. Jadhav VM, Thorat RM, Kadam VJ, Sathe NS. Hibiscus rosa-sinensis Linn — 'Rudrapuspa': a review. J Pharm Res. 2009 Jul;2(7):1168-73.
29. Semwal RB, Semwal DK, Combrinck S, Cartwright-Jones C, Viljoen A. Lawsonia inermis L. (henna): ethnobotanical, phytochemical and pharmacological aspects. J Ethnopharmacol. 2014 Aug 8;155(1):80-103. Available from: https://doi.org/10.1016/j.jep.2014.05.042
30. Gallo FR, Multari G, Palazzino G, Pagliuca G, Zappia G, Nicoletti M. Henna through the centuries: a quick HPTLC analysis proposal to check henna identity. Rev Bras Farmacogn. 2014 Mar-Apr;24(2):133-40. Available from: https://doi.org/10.1016/j.bjp.2014.03.008
31. Ponnusamy K, Petchiammal C, Mohankumar R, Hopper W. In vitro antifungal activity of indirubin isolated from a South Indian ethnomedicinal plant Wrightia tinctoria R. Br. J Ethnopharmacol. 2010 Oct 28;132(1):349-54. Available from:

https://doi.org/10.1016/j.jep.2010.07.050

32. Srivastava R. A review on phytochemical, pharmacological, and pharmacognostical profile of Wrightia tinctoria: an important medicinal plant. Pharmacogn Rev. 2014 Jan;8(15):36-44. Available from: https://doi.org/10.4103/0973-7847.125528
33. Gottumukkala VR, Annamalai T, Mukhopadhyay T. Phytochemical investigation and hair growth studies on the rhizomes of Nardostachys jatamansi DC. Pharmacogn Mag. 2011 Apr;7(26):146-50. Available from: https://doi.org/10.4103/0973-1296.80674
34. Purnima, Bhatt M, Kothiyal P. A review article on phytochemistry and pharmacological profiles of Nardostachys jatamansi DC. J Pharmacogn Phytochem. 2015;3(5):102-6.
35. Moldes AB, Rodríguez-López L, Rincón-Fontán M, López-Prieto A, Vecino X, Cruz JM. Synthetic and bio-derived surfactants versus microbial biosurfactants in the cosmetic industry: an overview. Int J Mol Sci. 2021 Feb 26;22(5):2371. Available from: https://doi.org/10.3390/ijms22052371
36. Katzbauer B. Properties and applications of xanthan gum. Polym Degrad Stab. 1998 Jan;59(1-3):81-4. Available from: https://doi.org/10.1016/S0141-3910(97)00180-8
37. Halla N, Fernandes IP, Heleno SA, Costa P, Boucherit-Otmani Z, Boucherit K, et al. Cosmetics preservation: a review on present strategies. Molecules. 2018 Jun 28;23(7):1571. Available from: https://doi.org/10.3390/molecules23071571
38. Bureau of Indian Standards (BIS). IS 7884:2023 / IS 7884:2004 Specification for Shampoo, Synthetic Detergent Based. New Delhi: Bureau of Indian Standards; 2023.
39. Ministry of AYUSH, Government of India. The Ayurvedic Pharmacopoeia of India (API), Part-II (Formulations), Volume-I to IV. New Delhi: Pharmacopoeia Commission for Indian Medicine & Homoeopathy; 2016.
40. Barnett G, Powers DH. A quantitative method for the evaluation of detergent action on hair. J Soc Cosmet Chem. 1951;2(3):219-28.
41. Luepke NP. Hen's egg chorioallantoic membrane test for irritation potential. Food Chem Toxicol. 1985 Feb;23(2):287-91. Available from: https://doi.org/10.1016/0278-6915(85)90030-4
42. Lademann J, Richter H, Teichmann A, Otberg N, Blume-Peytavi U, Luengo J, et al. Nanoparticles—an efficient carrier for drug delivery into the hair follicles. Eur J Pharm Biopharm. 2007 May;66(2):159-64. Available from: https://doi.org/10.1016/j.ejpb.2006.10.019
43. Patzelt A, Richter H, Knorr F, Schäfer U, Lehr CM, Dähne L, et al. Selective follicular targeting by modification of the particle sizes. J Control Release. 2011 Feb 28;150(1):45-8. Available from: https://doi.org/10.1016/j.jconrel.2010.11.015

44. Semalty A, Semalty M, Rawat BS, Singh D, Rawat MS. Pharmacosomes: the lipid-based new drug delivery system. Expert Opin Drug Deliv. 2009 Jun;6(6):599-612. Available from: https://doi.org/10.1517/17425240902967607
45. Lu M, Qiu Q, Luo X, Liu X, Sun J, Wang C, et al. Phyto-phospholipid complexes (phytosomes): a novel strategy to improve the bioavailability of active constituents. Asian J Pharm Sci. 2019 May;14(3):265-74. Available from: https://doi.org/10.1016/j.ajps.2018.05.011
46. Marianecci C, Di Marzio L, Rinaldi F, Celia C, Paolino D, Alhaique F, et al. Niosomes from 80s to present: the state of the art. Adv Colloid Interface Sci. 2014 Mar;205:187-206. Available from: https://doi.org/10.1016/j.cis.2013.11.018
47. Jantarat C, Sirathanarun P, Chuchue T, Konkamdee W, Ratanapongsai Y. Phyllanthus emblica extract-loaded transfersomes for hair follicle targeting: phytoconstituents, characterization, and hair growth promotion. J Oleo Sci. 2022;71(7):1085-96. Available from: https://doi.org/10.5650/jos.ess21425
48. Pardeike J, Hommoss A, Müller RH. Lipid nanoparticles (SLN, NLC) in cosmetic and pharmaceutical dermal products. Int J Pharm. 2009 Jan 21;366(1-2):170-84. Available from: https://doi.org/10.1016/j.ijpharm.2008.10.003
49. Vivek-Ananth RP, Mohanraj K, Sahoo AK, Samal A. IMPPAT 2.0: an enhanced and expanded phytochemical atlas of Indian medicinal plants. ACS Omega. 2023 Feb 24;8(9):8827-45. Available from: https://doi.org/10.1021/acsomega.3c00350
50. Hopkins AL. Network pharmacology: the next paradigm in drug discovery. Nat Chem Biol. 2008 Nov;4(11):682-90. Available from: https://doi.org/10.1038/nchembio.118
51. Adu SA, Naughton PJ, Marchant R, Banat IM. Microbial biosurfactants in cosmetic and personal skincare pharmaceutical formulations. Pharmaceutics. 2020 Nov 16;12(11):1099. Available from: https://doi.org/10.3390/pharmaceutics12111099